

Characteristics of EAS and Primary Particle Mass Composition in the Energy Region of $10^{17} - 3 \cdot 10^{19}$ eV by Yakutsk Data

S.P. Knurenko^a, V.P. Egorova^a, A.A. Ivanov^a, V.A. Kolosov^a, I.T. Makarov^a, Z.E. Petrov^a, I.Ye. Sleptsov^a, G.G. Struchkov^a.

^aYu.G.Shafer Institute of Cosmophysical Research and Aeronomy,
31 Lenin Ave., 677980 Yakutsk, Russia

In the work the data of the Yakutsk complex EAS array and their comparison with calculation in the case of primary nuclei of different chemical elements are presented. The calculation by QGSJET model have been used interpreting experimental data.

1. INTRODUCTION

It is not possible to measure a mass composition of primary cosmic rays (PCR) in the energy range $10^{15} - 10^{20}$ eV using the direct method. Nothing remains, but to resort to the indirect methods when for the similar estimates the measurement of different components of extensive air shower (EAS) are used. That can be characteristics of longitudinal or lateral development of shower in the air. And usually it is connected with the analysis of the most sensitive to the composition of shower components which differ from each other by the character of formatting and absorbing in the atmosphere, for example, of the charged particle flux (electron, muon), the flux of Čerenkov or ionization radiation.

2. ENERGY TRANSFERRED TO THE ELECTROMAGNETIC EAS COMPONENT

Fig.1 presents the EAS Yakutsk array experimental data and calculations by the model with the decelerated and moderate dissipation of the energy into the electromagnetic EAS component: quasiscaling models (solid line) and QGSJET (dashed line) [1]. From Fig.1 it is seen both the agreement of experimental data and calculations by QGSJET model (proton) in the region $E_0 \geq 3 \cdot 10^{18}$ eV, and disagreement at $E_0 \leq 3 \cdot 10^{18}$ eV. The scaling model gives a noticeably greater value




Figure 1. A portion of the energy transferred to the electromagnetic EAS component by Čerenkov light data at the Yakutsk array.

of E_m/E_0 in relation to the experimental data that is doubtlessly also connected with the break of scaling function in the region of ultra-high energies.

The experimental data in Fig.1 is well approximated by the expression of a form:

$$\frac{E_m}{E_0} = (0.964 \pm 0.011) - (0.079 \pm \pm 0.005) \cdot E_0^{-(0.147 \pm 0.008)}. \quad (1)$$

The relation (1) is primarily important for the comparison of estimations of E_0 obtained at the Yakutsk and Fly's Eye arrays [2].

The calculations in [2] (see Fig.1) have been carried out by the QGSJET model in the case of primary proton and iron nucleus. A good agreement of our calculations in the case of the primary proton is observed. The comparison of experimental data with calculations for the proton and iron nucleus indicates to the fact that the mass composition of particles of cosmic radiation in the energy region of $10^{17} - 10^{18}$ eV and above $3 \cdot 10^{18}$ eV must differ. At $E_0 \geq 3 \cdot 10^{18}$ eV the mass composition is most likely close to the proton one.

3. A PORTION OF MUONS WITH $E_{th} \geq 1$ GeV

The showers with $E_0 \geq 10^{18}$ eV and zenith angles $\theta < 60^\circ$ have been chosen. It is required so that the shower axis is within the array and not less than three muon detectors (one of them is at a distance of 1000 m from the shower axis) operate during the shower. The density of muons flux $\rho_\mu(1000)$ is like a median between the densities adjusted to $\langle R \rangle = 1000$ m, according to a mean lateral muon distribution function. In this case the detectors which have been operated in the shower (they have given zero indications), i.e. the detector threshold were used in analyze too. The muon flux densities at have distances of 300 and 600 m from the shower axis have been calculated in analogous way. Results are in Fig.2a and 2b. Calculations [3] have shown that $\rho_\mu(R)$ weekly depends on a zenith angle in the interval of $\Delta\theta = (0^\circ - 60^\circ)$. So for the comparison of calculations and experimental data we take a mean angle $\theta = 39^\circ$. In order to take into account both physical fluctuations in measurement of $\rho_\mu(1000)$ and methodical ones (apparatus errors and accuracy of the locate of the shower axis) we have considered the judicial interval in one $\sigma_\mu^{mean} = \sigma_\mu^{phys} + \sigma_\mu^{meth}$. In Fig.2a and 2b the calculation result is shown by dotted line in the case of primary proton and by a dashed line for the iron nuclei. A solid line shows the upper and lower limits for the case if EAS with $E_0 \geq 10^{19}$ eV would gener-




Figure 2. a) Portion of the muons with $E_{th} \geq 1$ GeV (%). $\rho_\mu(300)/\rho_s(300)$ (•) and $\rho_\mu(600)/\rho_s(600)$ (◦). b) $\rho_\mu(1000)$ vs. E_0 relations for observed events $10^{18} - 10^{19}$ eV (square), $10^{19} - 10^{20}$ eV (points) and $10^{20} - 10^{21}$ eV (triangles). Expected $\pm 1\sigma$ bounds for the distributions are indicated for proton, iron and gamma-ray primary by different curve as in the legend.

ate by a primary γ -quantum. The calculation for the primary γ -quantum has been taken from [4]. Dots for $\rho_\mu(300)$ and circles for $\rho_\mu(600)$ the experimental data show in Fig.2a. In Fig.2b the showers in the energy range of $10^{18} - 10^{19}$ eV are shown by squares, the showers with $E_0 \geq 10^{19}$ eV are shown by dots and the showers of maximum energy are shown by triangles. The comparison of experimental data presented in Fig.2a with calculations by the QGSJET model carried out for the case of the primary proton and iron nuclei confirms the hypothesis on the fact that the considerable portion of ultimate energy EAS have been formed by protons. Their portion decreases below the energy 10^{18} eV. It is seen from Fig.2b, the basic mass of points (showers with $E_0 \geq 10^{19}$ eV) falls into the interval for the proton. 23 points fall into a zone of superposition of a proton and iron nuclei and 19 showers from 116 fall into a zone of upper boundary of calculation for the primary γ -quantum. It testifies to the fact that even taking into account there exists a probability that showers of such energies can be generated by neutral particles and, in particular, by a primary γ -quantum. Then it is justified to use the analysis of directions of arrival of showers with $E_0 \geq 10^{19}$ eV for the search of the sources of highest energy cosmic rays. In this case the determination accuracy of arrival angles of such showers must be not worse than $(0.5^\circ - 1.5^\circ)$.

4. $X_{\max}$ FLUCTUATIONS

A large number of Čerenkov detectors operating in the individual EAS events and also the use of a new version of the QGSJET model allow to obtain quantitative estimations of mass composition of PCR. For this aim we have compared experimental data (see Fig.3) and theoretical predictions by the QGSJET model for different primary nucleus with the use of χ^2 criterion. The value of χ^2 has been determined by the equality

$$\chi^2(X_{\max}) = \sum \frac{[N_{\text{exp}}(X_{\max}) - N_{\text{theor}}(X_{\max})]^2}{N_{\text{theor}}(X_{\max})} \quad (2)$$

where $N_{\text{exp}}(X_{\max})$ is the experimental number of showers in the $\Delta X_{\max}$ interval. $N_{\text{theor}}(X_{\max}, A_i)$ is the analogous number of showers calculated un-




Figure 3. Depth of maximum distributions for difference fixes energy.

der the assumption that the mass number of the nucleus is equal to A_i , and $P(A_i)$ is the probability of the fact that the shower with the energy E_0 is formed by a primary particle A_i . Then:

$$N_{\text{theor}}(X_{\text{max}}) = \sum_{i=1}^n P(A_i) \cdot N_{\text{theor}}(X_{\text{max}}, A_i) \quad (3)$$

The analysis of form of the experimental distribution of X_{max} at optimal value of χ^2 with a definite portion of probability doesn't contradict to the following relationship for five nuclei component:

$$\bar{E}_0 = 5 \cdot 10^{17} \text{ eV: } p: (39 \pm 11)\%, \alpha: (31 \pm 13)\%, \\ M: (18 \pm 10)\%, H: (7 \pm 6)\%, Fe: (5 \pm 4)\%;$$

$$\bar{E}_0 = 1 \cdot 10^{18} \text{ eV: } p: (41 \pm 8)\%, \alpha: (32 \pm 11)\%, \\ M: (16 \pm 9)\%, H: (6 \pm 4)\%, Fe: (5 \pm 3)\% ;$$

$$\bar{E}_0 = 5 \cdot 10^{18} \text{ eV: } p: (60 \pm 14)\%, \alpha: (21 \pm 13)\%, \\ M: (10 \pm 8)\%, H: (5 \pm 4)\%, Fe: (3 \pm 3)\%.$$

Thus in the framework of the QGSJET model one can suppose that the mass composition of PCR changes transferring from the energy range $(5-30) \cdot 10^{17}$ eV to energy range $(5-30) \cdot 10^{18}$ eV. At $E_0 \geq 3 \cdot 10^{18}$ eV the primary cosmic radiation consists from $\sim 70\%$ protons and helium nuclei, a portion of the rest nuclei in the range where there is the second irregularity in the energetic spectrum type "ankle" doesn't exceed $\sim 30\%$. A high content of proton and helium nuclei in PCR in the region of formation of "ankle" is most likely connected with an appreciable contribution into the overall flux of cosmic radiation in the Earth's vicinity of radiation coming from beyond our Galaxy limits.

Acknowledgements

This work has been financially supported by RFBR, grant №02-02-16380, grant №03-02-17160 and grant INTAS №03-51-5112.

REFERENCES

1. Knurenko S.P. et al. // Proc. 26th ICRC. Salt Lake City. V.1. P.372.
2. Song C., Cao Z., Dawson B.R. et al. Astroparticle Phys. 14 (2000) 7-13.

3. Sleptsova V.R. et al. // Nucl. Phys. B (Proc. Suppl.). 2003.V.122. P. 255.
4. K. Shinozaki et al. // Proc. 28th ICRC. Tsukuba. Japan. 2003. V. 3. P. 401.